

**WESTERN NORTH CAROLINA  
HARD AND SOFT MAST  
SURVEY REPORT  
36<sup>th</sup> Year**

**FALL 2018**



**North Carolina Wildlife Resources Commission**

**Compiled and written by:  
Colleen Olfenbuttel  
Black bear and Furbearer Biologist  
November 2018**



© 2008 Arbor Day Foundation

Funding for the hard and soft mast survey was partially provided through a Pittman-Robertson Wildlife Restoration Grant. The Federal Aid in Wildlife Restoration Act, popularly known as the Pittman-Robertson Act, was approved by Congress on September 2, 1937, and began functioning July 1, 1938. The purpose of this Act was to provide funding for the selection, restoration, rehabilitation and improvement of wildlife habitat, wildlife management research, and the distribution of information produced by the projects. The Act was amended October 23, 1970, to include funding for hunter training programs and the development, operation and maintenance of public target ranges.

Funds are derived from an 11 percent Federal excise tax on sporting arms, ammunition, and archery equipment, and a 10 percent tax on handguns. These funds are collected from the manufacturers by the Department of the Treasury and are apportioned each year to the States and Territorial areas (except Puerto Rico) by the Department of the Interior on the basis of formulas set forth in the Act. Funds for hunter education and target ranges are derived from one-half of the tax on handguns and archery equipment.

Each state's apportionment is determined by a formula which considers the total area of the state and the number of licensed hunters in the state. The program is a cost-reimbursement program, where the state covers the full amount of an approved project then applies for reimbursement through Federal Aid for up to 75 percent of the project expenses. The state must provide at least 25 percent of the project costs from a non-federal source.



## **Introduction**

North Carolina Wildlife Resources Commission (NCWRC) personnel have surveyed hard mast in the Mountain Region of North Carolina since 1983. From 1983-2005, North Carolina's hard mast surveys were conducted and reported using a method developed by Whitehead (1969) with slight modifications (Wentworth et al. 1992). This same protocol was used in whole or part by Georgia and Tennessee for many years and was adopted by South Carolina in the 1990's. In an effort to reduce costs and manpower commitments, while maintaining quality data and standard methodology among neighboring states, the member states of the Southern Appalachian Black Bear Study Group (SABBSG, Georgia, North Carolina, South Carolina, and Tennessee) have long searched for an improved technique for monitoring hard mast surveys. Beginning with the 2006 survey, we are using a new protocol and formula for determining mast indices (Greenberg and Warburton 2007). The new protocol only requires simple calculation of percent crown with acorns in the field. In order to maintain consistency with the old technique, the new technique uses statistically verified equations to convert mast index values to numbers previously used with the Whitehead (1969) method. Hard mast results reported in this document utilize the techniques described in Greenberg and Warburton (2007) and are described using the scale used by our agency since 1983. Due to small sample sizes, results will no longer be reported for individual routes for hickory and beech, but overall values for these species will be reported. Sample sizes are sufficient to allow the reporting of values for both the white oak and red oak groups by route.

## **Hard Mast Overall Results**

The 2018 hard mast survey was conducted by WRC Land and Water Access staff, WRC Wildlife Management Division Private Lands staff, and South Mountains State Park staff on 12 routes in western North Carolina. A total of 1,394 trees were sampled including 539 from the white oak group, 682 from the red oak group, 138 hickories, and 35 beeches. Combining all groups of species, mast was rated as poor, with an overall index of 1.58, which is a decline from last year's mast crop index (3.44; Table 1). Since 1983, North Carolina has experienced 12 years out of 36 years in which the hard mast index was rated as poor. Including only the oak species, mast production rated as poor (1.61; Table 1).

White oak production rated as poor (0.94) and was below both the long-term average (1.86) below last year's index (2.13; Table 1). When the white oak group is separated by species, chestnut oak and white oak production rated as poor (0.62 and 1.32, respectively; Table 2). Red oak production rated as fair (2.14) and below the long-term average (2.83; Table 1) for the species. Separated by species, black oak and scarlet oak rated as fair and northern red oak rated as poor (3.99, 2.41, and 1.89, respectively; Table 2). Hickory production rated as poor (1.58) and below the long-term average (2.34) for the species (Table 1). Beech production (1.11) was poor and below the long-term average (4.04; Table 1).

## **Hard Mast Survey Area Results**

As in previous years, hard mast production varied by location and species (Table 3; Figures 1 and 2). However, there was more consistency among routes during in 2018. Ten areas surveyed had white oak productivity rated as poor, while the Standing Indian and Nantahala routes (Macon County) had fair to good white oak productivity, respectively (Table 3; Figure 1). Red oak production in 7 areas rated as poor, 2 areas rated as fair, and 3 areas rated as good (Table 3; Figure 2). Red oak productivity ranked highest at the lower and upper elevations and

was poorest between 3,000 and 4,900 feet (Table 4). White oak productivity also ranked highest at lower and upper elevations, but productivity was ranked poor across elevations gradients (Table 4).

### **Summer Soft Mast Survey Results**

A soft mast survey was implemented during the summer and fall of 1993 to document berry production and abundance. The technique used for evaluating the soft mast survey has remained consistent throughout this period including the current year. Summer soft mast surveys have been conducted in conjunction with the Sardine Bait Station Survey (SBSS). During summer 2006, based on an agreement with the member states of the SABBSG, we did not conduct the SBSS. Review of data from the SBSS indicates that we can obtain long-term bear population trend information by conducting the survey every other year. Because of the new schedule, the summer soft mast survey will be conducted in odd years. The previous survey was conducted in 2017 (Table 5 and 6) and the next survey was conducted during the summer of 2019.

### **Fall Soft Mast Survey Results**

The 2018 fall soft mast survey is conducted in conjunction with the hard mast survey. Overall, soft mast production was below the production observed in 2017; grape was slightly above the long-term average while pokeberry, cherry, and blackgum were below long-term averages (Table 7; Figure 3). Pokeberry, cherry and black gum rated as poor, while grape produced fair crops (Table 7). As observed in previous years, local areas experienced variable production of fall soft mast depending on species and area (Table 8).

### **Conclusion**

This season's hard mast crop was the twelfth year since 1983 in which the overall hard mast index was poor. The fall hard mast index was lower in 2018 than in 2017, and was consistently low across most locations, with most areas experiencing poor production. For example, several areas experienced very poor production of white oaks (Table 3; Figures 1 and 2). Red oak productivity (Fair; 2.14) was better than white oak productivity (Poor; 0.94; Table 1). Surrounding states, including Georgia, South Carolina and Virginia, also reported fair to poor hard mast production. Georgia reported poor white oak (1.4) and fair red oak (2.7) productivity. South Carolina also reported poor white oak (2.0) and fair red oak (3.4) and productivity. The overall trend in hard mast production shows a very slight declining trend since surveys were initiated in 1983 (Figure 4).

### **LITERATURE CITED**

- Greenberg, C.H., and G.S. Warburton. 2007. A fast and reliable hard mast index from acorn presence-absence tallies. *Journal of Wildlife Management* 71:1654-1661.
- Wentworth, J.M., A.S. Johnson, P.E. Hale, and K.E. Kammermeyer. 1992. Relationship of Acorn abundance and deer herd characteristics in the southern Appalachians. *Southern Journal of Applied Forestry* 16:5-8.
- Whitehead, C.J. 1969. Oak mast yields on wildlife management areas in Tennessee. Tennessee Game and Fish Commission, Nashville, USA.

Table 1. Hard Mast Survey Results for Western North Carolina, 1983-2018.

| <b>Year</b>    | <b>White Oak</b> | <b>Red Oak</b> | <b>All Oaks</b> | <b>Hickory</b> | <b>Beech</b> | <b>Total</b> |
|----------------|------------------|----------------|-----------------|----------------|--------------|--------------|
| 1983           | 1.43             | 2.59           |                 | 1.99           | 5.51         | 2.25         |
| 1984           | 1.08             | 2.73           |                 | 3.05           | 4.28         | 2.30         |
| 1985           | 2.01             | 3.66           |                 | 0.80           | 3.06         | 2.80         |
| 1986           | 1.32             | 1.98           |                 | 2.25           | 5.22         | 1.90         |
| 1987           | 1.16             | 0.56           |                 | 3.57           | 5.75         | 1.31         |
| 1988           | 3.16             | 4.07           |                 | 2.04           | 4.25         | 3.57         |
| 1989           | 0.43             | 4.89           |                 | 2.78           | 6.44         | 3.14         |
| 1990           | 1.85             | 2.62           |                 | 1.20           | 1.89         | 2.17         |
| 1991           | 2.38             | 1.93           |                 | 3.75           | 6.89         | 2.43         |
| 1992           | 1.07             | 2.45           |                 | 0.72           | 1.17         | 1.78         |
| 1993           | 0.65             | 3.58           |                 | 2.43           | 4.77         | 2.48         |
| 1994           | 2.06             | 3.48           |                 | 2.02           | 6.20         | 2.85         |
| 1995           | 2.80             | 5.60           |                 | 2.48           | 0.36         | 4.22         |
| 1996           | 3.70             | 1.99           |                 | 2.81           | 4.31         | 2.72         |
| 1997           | 0.53             | 1.79           |                 | 1.17           | 2.35         | 1.29         |
| 1998           | 2.26             | 4.68           |                 | 3.27           | 4.70         | 3.69         |
| 1999           | 3.28             | 2.76           |                 | 2.80           | 6.22         | 3.05         |
| 2000           | 0.50             | 2.11           |                 | 2.73           | 5.71         | 1.82         |
| 2001           | 2.83             | 4.92           |                 | 2.88           | 3.97         | 3.98         |
| 2002           | 1.90             | 3.01           |                 | 1.75           | 3.44         | 2.47         |
| 2003           | 1.24             | 0.68           |                 | 3.58           | 5.42         | 1.33         |
| 2004           | 3.99             | 2.93           |                 | 1.32           | 1.65         | 3.09         |
| 2005           | 0.70             | 3.11           |                 | 1.86           | 4.30         | 2.14         |
| 2006           | 1.70             | 1.40           | 1.50*           | 3.20           | 4.10         | 1.80         |
| 2007           | 3.02             | 1.19           | 2.04            | 0.73           | 2.71         | 1.90         |
| 2008           | 1.01             | 2.40           | 1.76            | 3.82           | 4.34         | 2.06         |
| 2009           | 0.48             | 2.47           | 1.55            | 1.72           | 5.58         | 1.67         |
| 2010           | 3.46             | 3.97           | 3.75            | 3.50           | 0.87         | 3.66         |
| 2011           | 1.17             | 2.22           | 1.74            | 1.30           | 4.96         | 1.76         |
| 2012           | 1.87             | 2.68           | 2.31            | 2.01           | 3.14         | 2.29         |
| 2013           | 1.00             | 1.43           | 1.23            | 2.43           | 4.45         | 1.44         |
| 2014           | 4.43             | 4.36           | 4.42            | 2.33           | 1.23         | 4.10         |
| 2015           | 1.07             | 2.65           | 1.92            | 2.64           | 5.77         | 2.09         |
| 2016           | 2.71             | 2.60           | 2.66            | 2.45           | 4.08         | 2.67         |
| 2017           | 2.13             | 4.42           | 3.40            | 3.20           | 5.69         | 3.44         |
| 2018           | 0.94             | 2.14           | 1.61            | 1.58           | 1.11         | 1.58         |
| <b>Average</b> | <b>1.86</b>      | <b>2.83</b>    | <b>2.30</b>     | <b>2.34</b>    | <b>4.05</b>  | <b>2.48</b>  |

Numerical Rating = Crop Quality

0.0 to 2.0 = Poor      2.1 to 4.0 = Fair

4.1 to 6.0 = Good      6.1 to 8.0 = Excellent

\* Not reported for prior years.

Table 2. Hard Mast Survey Results by Species, 2018.

| <b>Grouping</b> | <b>Species</b>              | <b>Index</b> | <b>Number of Trees Sampled</b> |
|-----------------|-----------------------------|--------------|--------------------------------|
| Hickories       | MH, PH, SH, GH <sup>1</sup> | 1.58         | 138                            |
| Beech           | Beech                       | 1.11         | 35                             |
| Red Oaks        | Black Oak                   | 3.99         | 30                             |
|                 | Northern Red Oak            | 1.89         | 471                            |
|                 | Scarlet Oak                 | 2.41         | 178                            |
| White Oaks      | Chestnut Oak                | 0.62         | 299                            |
|                 | White Oak                   | 1.32         | 240                            |

Numerical Rating = Crop Quality

0.0 to 2.0 = Poor      2.1 to 4.0 = Fair  
 4.1 to 6.0 = Good      6.1 to 8.0 = Excellent

<sup>1</sup>MH,SH, PH, GH: Mockernut Hickory, Pignut Hickory, Shagbark Hickory

Table 3. Hard Mast Survey Results by Area, 2018.

| <b>County</b>    | <b>Area</b>     | <b>White Oak</b> | <b>Red Oak</b> | <b>All Oaks</b> |
|------------------|-----------------|------------------|----------------|-----------------|
| Transylvania     | Avery Creek     | 0.6              | 1.3            | 1.0             |
| Haywood          | Cold Mountain   | 0.4              | 0.4            | 0.4             |
| Avery & Caldwell | Edgemont        | 0.4              | 1.1            | 0.8             |
| Clay             | Fires Creek     | 0.6              | 3.6            | 2.0             |
| Haywood          | Harmon Den      | 1.1              | 1.3            | 1.2             |
| Burke & McDowell | Linville Mtn.   | 0.6              | 1.0            | 0.7             |
| Macon            | Nantahala       | 5.5              | 4.4            | 4.7             |
| Mitchell         | Poplar          | 0.9              | 1.2            | 1.0             |
| Graham           | Santeetlah      | 0.7              | 4.4            | 2.8             |
| Haywood          | Sherwood        | 0.4              | 1.2            | 0.9             |
| Burke            | South Mountains | 1.8              | 2.7            | 2.2             |
| Macon            | Standing Indian | 3.4              | 4.8            | 4.2             |

Numerical Rating = Crop Quality

0.0 to 2.0 = Poor      2.1 to 4.0 = Fair  
 4.1 to 6.0 = Good      6.1 to 8.0 = Excellent




Figure 1. Red Oak Index by County in western North Carolina, 2018.




Figure 2. White Oak Index by County in western North Carolina, 2018.

Table 4. Hard Mast Survey Results by Elevation, 2018.

| <b>Elevation (ft.)</b> | <b>Red Oak</b> | <b>White Oak</b> |
|------------------------|----------------|------------------|
| <1900                  | 2.04           | 1.32             |
| 2000-2900              | 2.90           | 0.99             |
| 3000-3900              | 1.97           | 0.70             |
| 4000-4900              | 1.45           | 1.43             |
| 5000+                  | 4.31           | 1.36             |

---

Numerical Rating = Crop Quality

|                   |                        |
|-------------------|------------------------|
| 0.0 to 2.0 = Poor | 2.1 to 4.0 = Fair      |
| 4.1 to 6.0 = Good | 6.1 to 8.0 = Excellent |

Table 5. Results of Mountain Summer Soft Mast Surveys, 1993-2017<sup>1</sup>.

| <b>Year</b>    | <b>Blueberry</b> | <b>Huckleberry</b> | <b>Blackberry</b> | <b>Pokeberry</b> |
|----------------|------------------|--------------------|-------------------|------------------|
| 1993           | 3.24             | 3.56               | 3.81              | 2.44             |
| 1994           | 3.17             | 3.54               | 3.53              | 1.44             |
| 1995           | 1.92             | 2.46               | 3.12              | 1.20             |
| 1996           | 2.02             | 1.97               | 3.39              | 1.51             |
| 1997           | 2.84             | 2.95               | 3.78              | 1.96             |
| 1998           | 1.73             | 1.09               | 3.00              | 2.10             |
| 1999           | 2.72             | 2.45               | 2.90              | 1.78             |
| 2000           | 2.70             | 2.72               | 2.99              | 1.64             |
| 2001           | 2.27             | 2.73               | 2.87              | 0.87             |
| 2002           | 1.87             | 2.22               | 3.55              | 1.32             |
| 2003           | 2.27             | 2.74               | 3.20              | 1.02             |
| 2004           | 1.67             | 1.61               | 4.25              | 1.41             |
| 2005           | 1.57             | 1.41               | 4.07              | 1.48             |
| 2007           | 2.11             | 1.23               | 2.48              | 1.84             |
| 2009           | 2.08             | 2.06               | 2.78              | 1.09             |
| 2011           | 1.69             | 1.53               | 3.28              | 1.37             |
| 2013           | 1.87             | 1.07               | 3.73              | 1.89             |
| 2015           | 2.14             | 1.38               | 3.97              | 2.28             |
| 2017           | 1.64             | 1.15               | 2.74              | 1.04             |
| <b>Average</b> | <b>2.18</b>      | <b>2.10</b>        | <b>3.34</b>       | <b>1.56</b>      |

<sup>1</sup> After 2005, summer soft mast surveys are conducted every two years.

Table 6. Mountain Summer Soft Mast Survey Results by Area, 2017.

| <b>Area</b>            | <b>Blueberry</b> | <b>Huckleberry</b> | <b>Blackberry</b> | <b>Pokeberry</b> |
|------------------------|------------------|--------------------|-------------------|------------------|
| Daniel Boone           | 0.75             | 0.25               | 1.25              | 0.25             |
| Fires Creek/Santeetlah | 1.20             | 1.00               | 1.60              | 1.40             |
| Flattop                | 0.00             | 0.00               | 6.00              | 0.00             |
| Harmon Den Area        | 3.33             | 3.33               | 1.00              | 1.00             |
| Mt. Mitchell           | 1.33             | 0.67               | 2.00              | 0.33             |
| Pisgah Area            | 1.80             | 1.00               | 2.00              | 0.40             |
| Rich Mountain          | 1.50             | 1.00               | 3.00              | 1.00             |
| Standing Indian        | 0.00             | 0.43               | 0.29              | 0.00             |
| T. Chatham             | 2.25             | 1.75               | 1.75              | 0.75             |
| Cheoah                 | 1.00             | 1.00               | 1.00              | 1.50             |
| South Mountains        | 4.00             | 2.00               | 4.00              | 1.00             |
| Highlands              | 0.00             | 0.00               | 0.00              | 0.00             |
| Gorges State Park      | 6.00             | 3.00               | 4.00              | 2.00             |
| Lake James State Park  | 1.00             | 1.00               | 1.00              | 1.00             |
| Sandy Mush             | 1.00             | 2.00               | 9.00              | 6.00             |
| Green River            | 1.00             | 0.00               | 6.00              | 0.00             |
| <b>Average</b>         | <b>1.64</b>      | <b>1.15</b>        | <b>2.74</b>       | <b>1.04</b>      |

NA<sup>1</sup>=No summer soft mast data provided for this route.

Numerical Rating = Crop Quality

0.0 to 2.0 = Poor      2.1 to 4.0 = Fair

4.1 to 6.0 = Good      6.1 to 9.0 = Excellent

Table 7. Results of Mountain Fall Soft Mast Surveys, 1993-2018.

| <b>Year</b>    | <b>Pokeberry</b> | <b>Cherry</b> | <b>Grapes</b> | <b>Blackgum</b> |
|----------------|------------------|---------------|---------------|-----------------|
| 1993           | 2.00             | 2.71          | 2.14          | 0.43            |
| 1994           | 3.11             | 2.00          | 3.78          | 1.71            |
| 1995           | 2.67             | 5.00          | 2.22          | 1.78            |
| 1996           | 2.40             | 1.63          | 3.25          | 1.75            |
| 1997           | 4.20             | 1.25          | 3.14          | 0.75            |
| 1998           | 4.63             | 2.67          | 2.80          | 1.50            |
| 1999           | 2.40             | 2.70          | 3.25          | 1.10            |
| 2000           | 2.20             | 2.70          | 3.30          | 1.00            |
| 2001           | 2.80             | 3.30          | 4.18          | 2.33            |
| 2002           | 1.10             | 2.45          | 2.73          | 1.27            |
| 2003           | 2.33             | 3.00          | 2.55          | 2.22            |
| 2004           | 1.67             | 2.70          | 3.00          | 1.44            |
| 2005           | 2.45             | 2.09          | 1.36          | 1.55            |
| 2006           | 3.73             | 2.00          | 3.17          | 2.50            |
| 2007           | 2.08             | 1.58          | 2.73          | 0.67            |
| 2008           | 2.91             | 4.64          | 4.08          | 2.58            |
| 2009           | 1.92             | 1.82          | 2.33          | 1.83            |
| 2010           | 2.90             | 5.80          | 4.80          | 1.40            |
| 2011           | 2.50             | 1.67          | 2.33          | 1.42            |
| 2012           | 2.50             | 1.08          | 2.92          | 1.00            |
| 2013           | 2.00             | 2.75          | 2.75          | 1.08            |
| 2014           | 2.55             | 3.91          | 4.55          | 2.18            |
| 2015           | 2.17             | 2.09          | 2.23          | 1.82            |
| 2016           | 3.00             | 3.27          | 2.75          | 1.92            |
| 2017           | 2.73             | 1.82          | 2.45          | 1.18            |
| 2018           | 1.83             | 1.58          | 3.00          | 1.17            |
| <b>Average</b> | <b>2.57</b>      | <b>2.62</b>   | <b>2.99</b>   | <b>1.52</b>     |

Numerical Rating = Crop Quality

|                   |                        |
|-------------------|------------------------|
| 0.0 to 2.0 = Poor | 2.1 to 4.0 = Fair      |
| 4.1 to 6.0 = Good | 6.1 to 8.0 = Excellent |




Figure 3. Results of Mountain Fall Soft Mast Surveys by species, 1993-2018.

Table 8. Local Results of Mountain Fall Soft Mast Surveys, 2018.

| County           | Area            | Pokeberry   | Cherry      | Grapes      | Blackgum    |
|------------------|-----------------|-------------|-------------|-------------|-------------|
| Transylvania     | Avery Creek     | 2           | 2           | 4           | 2           |
| Haywood          | Cold Mountain   | 2           | 0           | 0           | 0           |
| Avery & Caldwell | Edgemont        | 2           | 0           | 4           | 2           |
| Clay             | Fires Creek     | 1           | 6           | 6           | 1           |
| Haywood          | Harmon Den      | 1           | 2           | 2           | 1           |
| Burke & McDowell | Linville Mtn.   | 4           | 2           | 6           | 2           |
| Macon            | Nantahala       | 0           | 0           | 0           | 0           |
| Mitchell         | Poplar          | 2           | 2           | 4           | 0           |
| Graham           | Santeetlah      | 2           | 2           | 4           | 1           |
| Haywood          | Sherwood        | 2           | 2           | 2           | 1           |
| Burke            | South Mountains | 4           | 1           | 4           | 4           |
| Macon            | Standing Indian | 0           | 0           | 0           | 0           |
| <b>Average:</b>  |                 | <b>1.83</b> | <b>1.58</b> | <b>3.00</b> | <b>1.17</b> |

Numerical Rating = Crop Quality

0.0 to 2.0 = Poor      2.1 to 4.0 = Fair

4.1 to 6.0 = Good      6.1 to 8.0 = Excellent




Figure 4. Annual hard mast index in western North Carolina, 1983 through 2018.